



# Bresenham's Line Algorithm

- Uses only incremental integer calculations
- Which pixel to draw ?
  - (11,11) or (11,12) ?
  - (51,50) or (51,49) ?
  - Answered by Bresenham



- For  $|m| < 1$ 
  - Start from left end point  $(x_0, y_0)$  step to each successive column ( $x$  samples) and plot the pixel whose scan line  $y$  value is closest to the line path.
  - After  $(x_k, y_k)$  the choice could be  $(x_k+1, y_k)$  or  $(x_k+1, y_k+1)$

$$y = m(x_k + 1) + b$$

Then


$$\begin{aligned}d_1 &= y - y_k \\ &= m(x_k + 1) + b - y_k\end{aligned}$$

And

$$\begin{aligned}d_2 &= (y_k + 1) - y \\ &= y_k + 1 - m(x_k + 1) - b\end{aligned}$$

Difference between separations

$$d_1 - d_2 = 2m(x_k + 1) - 2y_k + 2b - 1$$



## Defining decision parameter

$$p_k = \Delta x(d_1 - d_2) \quad [1]$$
$$= 2\Delta y \cdot x_k - 2\Delta x \cdot y_k + c$$

Sign of  $p_k$  is same as that of  $d_1 - d_2$  for  $\Delta x > 0$  (left to right sampling)

$$p_{k+1} = 2\Delta y \cdot x_{k+1} - 2\Delta x \cdot y_{k+1} + c$$

$$p_{k+1} - p_k = 2\Delta y(x_{k+1} - x_k) - 2\Delta x(y_{k+1} - y_k)$$

$$p_{k+1} = p_k + 2\Delta y - 2\Delta x(y_{k+1} - y_k)$$

For Recursive calculation, initially

$$p_0 = 2\Delta y - \Delta x$$

Constant =  $2\Delta y + \Delta x(2b - 1)$  Which is independent of pixel position

$c$  eliminated here

because  $x_{k+1} = x_k + 1$

$y_{k+1} - y_k = 0$  if  $p_k < 0$

$y_{k+1} - y_k = 1$  if  $p_k \geq 0$

Substitute  $b = y_0 - m \cdot x_0$   
and  $m = \Delta y / \Delta x$  in [1]

# Algorithm Steps ( $|m| < 1$ )

1. Input the two line endpoints and store the left endpoint in  $(x_0, y_0)$
2. Plot first point  $(x_0, y_0)$
3. Calculate constants  $\Delta x$ ,  $\Delta y$ ,  $2\Delta y$  and  $2\Delta y - 2\Delta x$ , and obtain  $p_0 = 2\Delta y - \Delta x$
4. At each  $x_k$  along the line, starting at  $k=0$ , perform the following test:  
If  $p_k < 0$ , the next point plot is  $(x_k+1, y_k)$  and
$$P_{k+1} = p_k + 2\Delta y$$
Otherwise, the next point to plot is  $(x_k + 1, y_k+1)$  and
$$P_{k+1} = p_k + 2\Delta y - 2\Delta x$$
5. Repeat step 4  $\Delta x$  times

# What's the advantage?

- Answer: involves only the calculation of constants  $\Delta x$ ,  $\Delta y$ ,  $2\Delta y$  and  $2\Delta y - 2\Delta x$  once and integer addition and subtraction in each steps